Author Index—Volume 38

A

Abbara, S.: *see* Ferencik, M.
Achenbach, S.: *see* Ferencik, M.; Regenfus, M.
Achenbach, S., Giesler, T., Ropers, D., Ulzheimer, S., Anders, K., Wenkel, E., Pohle, K., Kachelriess, M., Derlien, H., Kalender, W. A., Daniel, W. G., Bautz, W., Baum, U.: Comparison of image quality in contrast-enhanced coronary-artery visualization by electron beam tomography and retrospectively electrocardiogram-gated multislice spiral computed tomography, 119
Adam, G.: *see* Lorenzen, J.; Nolte-Ernsting, C.
Ahlström, H.: *see* Wikström, J.
Ahmad, T., Ohlsson, C., Östenson, C.-G., Kreicbergs, A.: Peripheral quantitative computed tomography for the detection of diabetic osteopathy: A study in the Goto-Kakizaki rat, 171
Alfke, H.: *see* Kalinowski, M.
Alves, F. C., Donato, P., Sherry, A. D., Zaheer, A., Zhang, S., Lubag, A. J. M., Merritt, M. E., Lenkinski, R. E., Frangioni, J. V., Neves, M., Prata, M. I. M., Santos, A. C., de Lima, J. J. P., Geraldes, C. F. G. C.: Silencing of phosphonate-gadolinium magnetic resonance imaging contrast by hydroxyapatite binding, 750
Anders, K.: *see* Achenbach, S.
Angele, P.: *see* Völk, M.
Antoch, G., Jentzen, W., Freudenberg, L. S., Stattaus, J., Mueller, S. P., Debatin, J. F., Bockisch, A.: Effect of oral contrast agents on computed tomography-based positron emission tomography attenuation correction in dual-modality positron emission tomography/computed tomography imaging, 784
Aubé, C.: *see* Brieger, J.
Averbukh, M.: *see* Peer, A.
Averbukh, Z.: *see* Peer, A.
Azhari, T.: *see* Pfleiderer, S. O. R.

B

Baik, S. K.: *see* Cha, S.-H.
Barentsz, J. O.: *see* Wikström, J.
Barth, M., Nöbauer-Huhmann, I.-M., Reichenbach, J. R., Mlynárik, V., Schöggl, A., Matula, C., Trattnig, S.: High-resolution three-dimensional contrast-enhanced blood oxygenation level-dependent magnetic resonance venography of brain tumors at 3 Tesla: First clinical experience and comparison with 1.5 Tesla, 409
Ba-Ssalamah, A., Nöbauer-Huhmann, I. M., Pinker, K., Schibany, N., Prokesch, R., Mehrain, S., Mlynárik, V., Fog, A., Heimberger, K., Trattnig, S.: Effect of contrast dose and field strength in the magnetic resonance detection of brain metastases, 415
Baum, U.: *see* Achenbach, S.
Bautz, W.: *see* Achenbach, S.
Becker, C. R., Hong, C., Knez, A., Leber, A., Bruening, R., Schoepf, U. J., Reiser, M. F.: Optimal contrast application for cardiac 4-detector-row computed tomography, 690
Belenger, J.: *see* Montet, X.
Berg, A.: *see* Langenberger, H.
Berg, A., Sailer, J., Rand, T., Moser, E.: Diffusivity- and T2 imaging at 3 Tesla for the detection of degenerative changes in human-excised tissue with high resolution: Atherosclerotic arteries, 452
Berg, A., Singer, T., Moser, E.: High-resolution diffusivity imaging at 3.0 T for the detection of degenerative changes: A trypsin-based arthritis model, 460
Berman, S.: *see* Peer, A.
Berthezène, Y.: *see* Stupar, V.
Bick, U.: *see* Lawaczeck, R.
Biesterfeldt, M.: *see* Lorenzen, J.
Birkfellner, W.: *see* Wagner, A.
Blackband, S. J.: *see* Kim, H.-W.
Blackshear, J. L.: *see* Gerber, T. C.
Bley, T. A.: *see* Kotter, E.
Bloebaum, R. D.: *see* Kuo, T. Y.
Bock, M.: *see* Fink, C.
Bockisch, A.: *see* Antoch, G.
Bocksch, W.: *see* Knollmann, F. D.
Boehm, H. F., Raeth, C., Monetti, R. A., Mueller, D., Newitt, D., Majumdar, S., Rummeny, E., Morfill, G., Link, T. M.: Local 3D scaling properties for the analysis of trabecular bone extracted from high-resolution magnetic resonance imaging of human trabecular bone: Comparison with bone mineral density in the prediction of biomechanical strength in vitro, 269
Boesinger, P.: *see* Spiegel, M. A.
Böhm, I., Speck, U., Schild, H.: Cytokine profiles after nonionic dimeric contrast medium injection, 776
Böhm, P.: *see* König, C. W.
Bongers, A.: *see* Weber, M.-A.
Bonomo, L.: *see* Wikström, J.
Booth, S. L.: *see* Ferencik, M.
Börnert, P.: *see* Spuentrup, E.
Bosk, S.: *see* Herborn, C. U.
Botnar, R.: *see* Spuentrup, E.
Botnar, R. M.: *see* Spuentrup, E.
Bouvier, S.: *see* Sicotte, N. L.
Box, F. M. A., Spilt, A., Van Buchem, M. A., van der Geest, R. J., Reiber, J. H. C.: Automatic model-based contour detection and blood flow quantification in small vessels with velocity encoded magnetic resonance imaging, 567
Brady, T. J.: *see* Ferencik, M.
Bräuer, R.: *see* Hansch, A.
Brechbiel, M.: *see* Wang, S. J.
Breithardt, G.: *see* Grude, M.
Brieger, J., Pereira, P. L., Trübenbach, J., Schenk, M., Kröber, S.-M., Schmidt, D., Aubé, C., Claussen, C. D., Schick, F.: In vivo efficiency of four commercial monopolar radiofrequency ablation systems: A comparative experimental study in pig liver, 609
Briguet, A.: *see* Chaabane, L.
Bruening, R.: *see* Becker, C. R.
Buckley, D. L.: *see* Kim, H.-W.
Budoff, M. J.: *see* Mao, S.
Buecker, A.: *see* Spuentrup, E.
Bui, C. D., Ching, A. S., Carlos, R. C., Shreve, P. D., Mukherji, S. K.: Diagnostic accuracy of 2-[fluorine-18]fluro-2-deoxy-D-glucose positron emission tomography imaging in nonsquamous tumors of the head and neck, 593
Burnett, R.: *see* Labarthe, B.
Burtea, C., Laurent, S., Colet, J.-M., Vander Elst, L. V., Muller, R. N.: Development of new glucosylated derivatives of gadolinium diethylenetriaminepentaacetic for magnetic resonance angiography, 320
Busch, M.: *see* Kivelitz, D.

C

Canet, E.: *see* Stupar, V.
Carlos, R. C.: *see* Bui, C. D.
Carson, S.: *see* Mao, S.
Caserta, J.: *see* Kim, H.-W.; Peterson, D. M.
Cha, S.-H., Han, M. H., Choi, Y. H., Yoon, C. J., Baik, S. K., Kim, S. J., Chang, K.-H.: Vascular responses in normal canine carotid arteries: Comparison between various self-expanding stents of the same unconstrained size, 95
Chaabane, L., Soulas, E. C., Contard, F., Salah, A., Guerrier, D., Briguet, A., Douek, P.: High-resolution magnetic resonance imaging at 2 Tesla: Potential for atherosclerotic lesions exploration in the apolipoprotein E knockout mouse, 532
Chan, D. Y.: *see* Solomon, S. B.
Chan, R. C.: *see* Ferencik, M.
Chang, K.-H.: *see* Cha, S.-H.; Kim, Y. H.; Kwon, O.-K.
Chapon, C., Franconi, F., Lemaire, L., Marescaux, L., Legras, P., Saint-André, J. P., Denizot, B., Le Jeune, J.-J.: High field magnetic resonance imaging evaluation of superparamagnetic iron oxide nanoparticles in a permanent rat myocardial infarction, 141
Cheon, J.-E., Kim, I.-O., Kim, C. J., Kim, W. S., Yoo, W. J., Choi, I. H., Yeon, K. M.: Imaging findings after fat graft interposition in an injured growth plate: An experimental study in rabbits, 695
Child, J.: *see* Mao, S.
Ching, A. S.: *see* Bui, C. D.
Cho, W. H.: *see* Kim, Y. H.
Choe, G.: *see* Kwon, O.-K.
Choi, B. I.: *see* Kim, Y. H.
Choi, I. H.: *see* Cheon, J.-E.
Choi, Y. H.: *see* Cha, S.-H.
Chung, G. H.: *see* Lee, J. M.
Chung, M. J.: *see* Lee, J. M.
Chung, S. K.: *see* Kim, K.
Chung, Y. A.: *see* Kim, K.
Clauβ, W., Scheller, B., Schmitt, A., Sovak, M., Speck, U.: No difference among modern contrast media's effect on neointimal proliferation and restenosis after coronary stenting in pigs, 743
Claussen, C. D.: *see* Brieger, J.
Cohen, M. S.: *see* Sicotte, N. L.
Colet, J.-M.: *see* Burtea, C.
Contard, F.: *see* Chaabane, L.
Corot, C.: *see* Labarthe, B.
Corot, C., Violas, X., Robert, P., Gagneur, G., Port, M.: Comparison of different types of blood pool agents (P792, MS325, USPIO) in a rabbit MR angiography-like protocol, 311
Crémillieux, Y.: *see* Stupar, V.
Crisp, D. E.: *see* Desai, N.
Crisp, M. B.: *see* Desai, N.

D

Damilakis, J., Papadokostakis, G., Vrahoriti, H., Tsagaraki, I., Perisinakis, K., Hadjipavlou, A., Gourtsoyiannis, N.: Ultrasound velocity through the cortex of phalanges, radius, and tibia in normal and osteoporotic postmenopausal women using a new multisite quantitative ultrasound device, 207
Danias, P. G.: *see* Storey, P.
Daniel, W. G.: *see* Achenbach, S.; Regenfus, M.
Daprà, M.: *see* Wikström, J.
Debatin, J. F.: *see* Antoch, G.; Herborn, C. U.

Author Index—Volume 38

A

Abbara, S.: *see* Ferencik, M.
Achenbach, S.: *see* Ferencik, M.; Regenfus, M.
Achenbach, S., Giesler, T., Ropers, D., Ulzheimer, S., Anders, K., Wenkel, E., Pohle, K., Kachelriess, M., Derlien, H., Kalender, W. A., Daniel, W. G., Bautz, W., Baum, U.: Comparison of image quality in contrast-enhanced coronary-artery visualization by electron beam tomography and retrospectively electrocardiogram-gated multislice spiral computed tomography, 119
Adam, G.: *see* Lorenzen, J.; Nolte-Ernsting, C.
Ahlström, H.: *see* Wikström, J.
Ahmad, T., Ohlsson, C., Östenson, C.-G., Kreicbergs, A.: Peripheral quantitative computed tomography for the detection of diabetic osteopathy: A study in the Goto-Kakizaki rat, 171
Alfke, H.: *see* Kalinowski, M.
Alves, F. C., Donato, P., Sherry, A. D., Zaheer, A., Zhang, S., Lubag, A. J. M., Merritt, M. E., Lenkinski, R. E., Frangioni, J. V., Neves, M., Prata, M. I. M., Santos, A. C., de Lima, J. J. P., Geraldes, C. F. G. C.: Silencing of phosphonate-gadolinium magnetic resonance imaging contrast by hydroxyapatite binding, 750
Anders, K.: *see* Achenbach, S.
Angele, P.: *see* Völk, M.
Antoch, G., Jentzen, W., Freudenberg, L. S., Stattaus, J., Mueller, S. P., Debatin, J. F., Bockisch, A.: Effect of oral contrast agents on computed tomography-based positron emission tomography attenuation correction in dual-modality positron emission tomography/computed tomography imaging, 784
Aubé, C.: *see* Brieger, J.
Averbukh, M.: *see* Peer, A.
Averbukh, Z.: *see* Peer, A.
Azhari, T.: *see* Pfleiderer, S. O. R.

B

Baik, S. K.: *see* Cha, S.-H.
Barentsz, J. O.: *see* Wikström, J.
Barth, M., Nöbauer-Huhmann, I.-M., Reichenbach, J. R., Mlynárik, V., Schöggl, A., Matula, C., Trattnig, S.: High-resolution three-dimensional contrast-enhanced blood oxygenation level-dependent magnetic resonance venography of brain tumors at 3 Tesla: First clinical experience and comparison with 1.5 Tesla, 409
Ba-Ssalamah, A., Nöbauer-Huhmann, I. M., Pinker, K., Schibany, N., Prokesch, R., Mehrain, S., Mlynárik, V., Fog, A., Heimberger, K., Trattnig, S.: Effect of contrast dose and field strength in the magnetic resonance detection of brain metastases, 415
Baum, U.: *see* Achenbach, S.
Bautz, W.: *see* Achenbach, S.
Becker, C. R., Hong, C., Knez, A., Leber, A., Bruening, R., Schoepf, U. J., Reiser, M. F.: Optimal contrast application for cardiac 4-detector-row computed tomography, 690
Belenger, J.: *see* Montet, X.
Berg, A.: *see* Langenberger, H.
Berg, A., Sailer, J., Rand, T., Moser, E.: Diffusivity- and T2 imaging at 3 Tesla for the detection of degenerative changes in human-excised tissue with high resolution: Atherosclerotic arteries, 452
Berg, A., Singer, T., Moser, E.: High-resolution diffusivity imaging at 3.0 T for the detection of degenerative changes: A trypsin-based arthritis model, 460
Berman, S.: *see* Peer, A.
Berthezène, Y.: *see* Stupar, V.
Bick, U.: *see* Lawaczeck, R.
Biesterfeldt, M.: *see* Lorenzen, J.
Birkfellner, W.: *see* Wagner, A.
Blackband, S. J.: *see* Kim, H.-W.
Blackshear, J. L.: *see* Gerber, T. C.
Bley, T. A.: *see* Kotter, E.
Bloebaum, R. D.: *see* Kuo, T. Y.
Bock, M.: *see* Fink, C.
Bockisch, A.: *see* Antoch, G.
Bocksch, W.: *see* Knollmann, F. D.
Boehm, H. F., Raeth, C., Monetti, R. A., Mueller, D., Newitt, D., Majumdar, S., Rummeny, E., Morfill, G., Link, T. M.: Local 3D scaling properties for the analysis of trabecular bone extracted from high-resolution magnetic resonance imaging of human trabecular bone: Comparison with bone mineral density in the prediction of biomechanical strength in vitro, 269
Boesinger, P.: *see* Spiegel, M. A.
Böhm, I., Speck, U., Schild, H.: Cytokine profiles after nonionic dimeric contrast medium injection, 776
Böhm, P.: *see* König, C. W.
Bongers, A.: *see* Weber, M.-A.
Bonomo, L.: *see* Wikström, J.
Booth, S. L.: *see* Ferencik, M.
Börnert, P.: *see* Spuentrup, E.
Bosk, S.: *see* Herborn, C. U.
Botnar, R.: *see* Spuentrup, E.
Botnar, R. M.: *see* Spuentrup, E.
Bouvier, S.: *see* Sicotte, N. L.
Box, F. M. A., Spilt, A., Van Buchem, M. A., van der Geest, R. J., Reiber, J. H. C.: Automatic model-based contour detection and blood flow quantification in small vessels with velocity encoded magnetic resonance imaging, 567
Brady, T. J.: *see* Ferencik, M.
Bräuer, R.: *see* Hansch, A.
Brechbiel, M.: *see* Wang, S. J.
Breithardt, G.: *see* Grude, M.
Brieger, J., Pereira, P. L., Trübenbach, J., Schenk, M., Kröber, S.-M., Schmidt, D., Aubé, C., Claussen, C. D., Schick, F.: In vivo efficiency of four commercial monopolar radiofrequency ablation systems: A comparative experimental study in pig liver, 609
Briguet, A.: *see* Chaabane, L.
Bruening, R.: *see* Becker, C. R.
Buckley, D. L.: *see* Kim, H.-W.
Budoff, M. J.: *see* Mao, S.
Buecker, A.: *see* Spuentrup, E.
Bui, C. D., Ching, A. S., Carlos, R. C., Shreve, P. D., Mukherji, S. K.: Diagnostic accuracy of 2-[fluorine-18]fluro-2-deoxy-D-glucose positron emission tomography imaging in nonsquamous tumors of the head and neck, 593
Burnett, R.: *see* Labarthe, B.
Burtea, C., Laurent, S., Colet, J.-M., Vander Elst, L. V., Muller, R. N.: Development of new glucosylated derivatives of gadolinium diethylenetriaminepentaacetic for magnetic resonance angiography, 320
Busch, M.: *see* Kivelitz, D.

C

Canet, E.: *see* Stupar, V.
Carlos, R. C.: *see* Bui, C. D.
Carson, S.: *see* Mao, S.
Caserta, J.: *see* Kim, H.-W.; Peterson, D. M.
Cha, S.-H., Han, M. H., Choi, Y. H., Yoon, C. J., Baik, S. K., Kim, S. J., Chang, K.-H.: Vascular responses in normal canine carotid arteries: Comparison between various self-expanding stents of the same unconstrained size, 95
Chaabane, L., Soulas, E. C., Contard, F., Salah, A., Guerrier, D., Briguet, A., Douek, P.: High-resolution magnetic resonance imaging at 2 Tesla: Potential for atherosclerotic lesions exploration in the apolipoprotein E knockout mouse, 532
Chan, D. Y.: *see* Solomon, S. B.
Chan, R. C.: *see* Ferencik, M.
Chang, K.-H.: *see* Cha, S.-H.; Kim, Y. H.; Kwon, O.-K.
Chapon, C., Franconi, F., Lemaire, L., Marescaux, L., Legras, P., Saint-André, J. P., Denizot, B., Le Jeune, J.-J.: High field magnetic resonance imaging evaluation of superparamagnetic iron oxide nanoparticles in a permanent rat myocardial infarction, 141
Cheon, J.-E., Kim, I.-O., Kim, C. J., Kim, W. S., Yoo, W. J., Choi, I. H., Yeon, K. M.: Imaging findings after fat graft interposition in an injured growth plate: An experimental study in rabbits, 695
Child, J.: *see* Mao, S.
Ching, A. S.: *see* Bui, C. D.
Cho, W. H.: *see* Kim, Y. H.
Choe, G.: *see* Kwon, O.-K.
Choi, B. I.: *see* Kim, Y. H.
Choi, I. H.: *see* Cheon, J.-E.
Choi, Y. H.: *see* Cha, S.-H.
Chung, G. H.: *see* Lee, J. M.
Chung, M. J.: *see* Lee, J. M.
Chung, S. K.: *see* Kim, K.
Chung, Y. A.: *see* Kim, K.
Clauβ, W., Scheller, B., Schmitt, A., Sovak, M., Speck, U.: No difference among modern contrast media's effect on neointimal proliferation and restenosis after coronary stenting in pigs, 743
Claussen, C. D.: *see* Brieger, J.
Cohen, M. S.: *see* Sicotte, N. L.
Colet, J.-M.: *see* Burtea, C.
Contard, F.: *see* Chaabane, L.
Corot, C.: *see* Labarthe, B.
Corot, C., Violas, X., Robert, P., Gagneur, G., Port, M.: Comparison of different types of blood pool agents (P792, MS325, USPIO) in a rabbit MR angiography-like protocol, 311
Crémillieux, Y.: *see* Stupar, V.
Crisp, D. E.: *see* Desai, N.
Crisp, M. B.: *see* Desai, N.

D

Damilakis, J., Papadokostakis, G., Vrahoriti, H., Tsagaraki, I., Perisinakis, K., Hadjipavlou, A., Gourtsoyiannis, N.: Ultrasound velocity through the cortex of phalanges, radius, and tibia in normal and osteoporotic postmenopausal women using a new multisite quantitative ultrasound device, 207
Danias, P. G.: *see* Storey, P.
Daniel, W. G.: *see* Achenbach, S.; Regenfus, M.
Daprà, M.: *see* Wikström, J.
Debatin, J. F.: *see* Antoch, G.; Herborn, C. U.

Debus, J.: *see* Weber, M.-A.
de Lima, J. J. P.: *see* Alves, F. C.
Del Maschio, A.: *see* Wikström, J.
Delorme, S.: *see* Fink, C.; Kiessling, F.; Weber, M.-A.
DeMaria, A. N.: *see* Masugata, H.
Dembinski, R.: *see* Wildberger, J. E.
Denizot, B.: *see* Chapon, C.
Derlien, H.: *see* Achenbach, S.
Desai, N., Runge, V. M., Crisp, D. E., Crisp, M. B., Naul, L. G.: Magnetic resonance imaging of the brain in glutaric acidemia type I: A review of the literature and a report of four new cases with attention to the basal ganglia and imaging technique, 489
Deutsch, J.: *see* Grenacher, L.
Dickhoff, P.: *see* Frayne, R.
Diekmann, F.: *see* Lawaczeck, R.
Diekmann, S.: *see* Lawaczeck, R.
Djurhuus, J. C.: *see* Rawashdeh, Y. F.
Donato, P.: *see* Alves, F. C.
Douek, P.: *see* Chaabane, L.
Duda, S. H.: *see* König, C. W.
Duensing, G. R.: *see* Kim, H.-W.; Peterson, D. M.
Dupuich, D.: *see* Stupar, V.

E

Eberle, B.: *see* Gast, K. K.
Edelman, R. R.: *see* Storey, P.
Einert, A.: *see* Kotter, E.
Endoh, M.: *see* Tsuda, M.
Essig, M.: *see* Weber, M.-A.
Ewers, R.: *see* Wagner, A.

F

Fallenberg, E. M.: *see* Grude, M.
Farhan, N.: *see* Kiessling, F.
Federle, M. P.: *see* Murakami, T.
Felix, R.: *see* Knollmann, F. D.
Ferencik, M., Ferullo, A., Achenbach, S., Abbara, S., Chan, R. C., Booth, S. L., Brady, T. J., Hoffmann, U.: Coronary calcium quantification using various calibration phantoms and scoring thresholds, 559
Ferlitsch, K.: *see* Langenberger, H.
Ferrara, K. W.: *see* Wisner, E. R.
Ferullo, A.: *see* Ferencik, M.
Feuerbach, S.: *see* Hamer, O. W.; Völk, M.
Fink, C.: *see* Kiessling, F.
Fink, C., Bock, M., Puderbach, M., Schmähl, A., Delorme, S.: Partially parallel three-dimensional magnetic resonance imaging for the assessment of lung perfusion–Initial results, 482
Fisch, D.: *see* Kotter, E.
Fischbach, R.: *see* Grude, M.; Maintz, D.
Fitzsimmons, J.: *see* Kim, H.-W.
Fitzsimmons, J. R.: *see* Peterson, D. M.
Fjield, T.: *see* Solomon, S. B.
Flohr, T.: *see* Maintz, D.
Flohr, T. G.: *see* Mahnken, A. H.
Fog, A.: *see* Ba-Ssalamah, A.
Franconi, F.: *see* Chapon, C.
Frangioni, J. V.: *see* Alves, F. C.
Frayne, R., Goodyear, B. G., Dickhoff, P., Lauzon, M. L., Sevick, R. J.: Magnetic resonance imaging at 3.0 Tesla: Challenges and advantages in clinical neurological imaging, 385
Freudenberg, L. S.: *see* Antoch, G.
Frey, O.: *see* Hansch, A.
Frey, R.: *see* Gruber, S.
Fried, N.: *see* Solomon, S. B.
Fritz, J.: *see* König, C. W.
Frokiaer, J.: *see* Rawashdeh, Y. F.
Fuchsjäger, M.: *see* Malich, A.
Füchtmeier, B.: *see* Völk, M.
Fusenig, N. E.: *see* Kiessling, F.

G

Gabe, J. D.: *see* Wisner, E. R.
Gagneur, G.: *see* Corot, C.
Gallowitsch, H.-J., Kresnik, E., Gasser, J., Kumnig, G., Igerc, I., Mikosch, P., Lind, P.: F-18 fluorodeoxyglucose positron-emission tomography in the diagnosis of tumor recurrence and metastases in the follow-up of patients with breast carcinoma: A comparison to conventional imaging, 250
Gasser, J.: *see* Gallowitsch, H.-J.
Gast, K. K., Puderbach, M. U., Rodriguez, I., Eberle, B., Markstaller, K., Knitz, F., Schmiedeskamp, J., Weiler, N., Schreiber, W. G., Mayer, E., Thelen, M., Kauczor, H.-U.: Distribution of ventilation in lung transplant recipients: Evaluation by dynamic ^{3}He-MRI with lung motion correction, 341
Gehl, H.-B.: *see* Schneider, G.
Geraldes, C. F. G. C.: *see* Alves, F. C.
Gerber, T. C., O'Brien, P. C., Pastor, K., Kuzo, R. S., Blackshear, J. L., Morin, R. L.: Evaluation of reconstruction windows for multislice computed tomography in quantification of coronary calcium, 108
Ghanem, N.: *see* Kotter, E.
Giesler, T.: *see* Achenbach, S.
Gilles, T.: *see* Krause, W.
Glowinski, A.: *see* Nolte-Ernsting, C.
Goodyear, B. G.: *see* Frayne, R.
Gourtsoyiannis, N.: *see* Damilakis, J.
Goyen, M.: *see* Herborn, C. U.
Grampp, S.: *see* Langenberger, H.; Wagner, A.
Grenacher, L., Deutsch, J., Lubienski, A., Richter, G. M.: Resistance to hoop stress in balloon expandable stents: Evaluation in an ex vivo model, 65
Gruber, S., Frey, R., Mlynárik, V., Stadlbauer, A., Heiden, A., Kasper, S., Kemp, G. J., Moser, E.: Quantification of metabolic differences in the frontal brain of depressive patients and controls obtained by ^{1}H-MRS at 3 Tesla, 403
Grude, M., Juergens, K. U., Wichter, T., Paul, M., Fallenberg, E. M., Muller, J. G., Heindel, W., Breithardt, G., Fischbach, R.: Evaluation of global left ventricular myocardial function with electrocardiogram-gated multidetector computed tomography: Comparison with magnetic resonance imaging, 653
Gudinchet, F.: *see* Meuwly, J.-Y.
Guerrier, D.: *see* Chaabane, L.
Günther, M.: *see* Weber, M.-A.
Günther, R. W.: *see* Mahnken, A. H.; Nolte-Ernsting, C.; Spuentrup, E.; Wildberger, J. E.

H

Haas, M.: *see* Hansch, A.
Hadjipavlou, A.: *see* Damilakis, J.
Hamer, O.: *see* Völk, M.
Hamer, O. W., Völk, M., Zorger, N., Feuerbach, S., Strotzer, M.: Amorphous silicon, flat-panel, x-ray detector *versus* storage phosphor-based computed radiography: Contrast-detail phantom study at different tube voltages and detector entrance doses, 212
Hamm, B.: *see* Kivelitz, D.; Lawaczeck, R.; Wikström, J.
Han, D. H.: *see* Kwon, O.-K.
Han, J. K.: *see* Kim, Y. H.
Han, M. H.: *see* Cha, S.-H.; Kwon, O.-K.
Han, Y. M.: *see* Lee, J. M.
Hansch, A., Sauner, D., Hilger, I., Frey, O., Haas, M., Malich, A., Bräuer, R., Kaiser, W. A.: Noninvasive diagnosis of arthritis by autofluorescence, 578
Harnish, P. P.: *see* Storey, P.
Hazle, J. D.: *see* Kangasniemi, M.
He, X.: *see* Shin, J. H.
Heiden, A.: *see* Gruber, S.
Heilmann, M.: *see* Kiessling, F.
Heimberger, K.: *see* Ba-Ssalamah, A.
Heindel, W.: *see* Grude, M.; Maintz, D.
Helmig, K.: *see* Knollmann, F. D.
Hente, R.: *see* Völk, M.
Hentrich, H.-R.: *see* Wikström, J.
Herborn, C. U., Lauenstein, T. C., Ruehm, S. G., Bosk, S., Debatin, J. F., Goyen, M.: Intraindividual comparison of gadopentetate dimeglumine, gadobenate dimeglumine, and gadobutrol for pelvic 3D magnetic resonance angiography, 27
Hetzer, R.: *see* Knollmann, F. D.
Heussel, C. P., Scholz, A., Schmittner, M., Laukemper-Ostendorf, S., Schreiber, W. G., Ley, S., Quintel, M., Weiler, N., Thelen, M., Kauczor, H.-U.: Measurements of alveolar pO_2 using ^{19}F-MRI in partial liquid ventilation, 635
Heussen, N.: *see* Mahnken, A. H.
Hilger, I.: *see* Hansch, A.
Hinton, D. P., Wald, L. L., Pitts, J., Schmitt, F.: Comparison of cardiac MRI on 1.5 and 3.0 Tesla clinical whole body systems, 436
Hoffmann, U.: *see* Ferencik, M.
Hong, C.: *see* Becker, C. R.
Hori, M.: *see* Murakami, T.
Hørlyck, A.: *see* Rawashdeh, Y. F.
Hummel, M.: *see* Knollmann, F. D.
Hvistendahl, J. J.: *see* Rawashdeh, Y. F.

I

Idee, J.-M.: *see* Labarthe, B.
Igerc, I.: *see* Gallowitsch, H.-J.
Ishibashi, T.: *see* Tsuda, M.
Ito, Y.: *see* Sumimura, T.
Itoh, Y.: *see* Sumimura, T.
Ivancevic, M. K.: *see* Montet, X.

J

Jackson, E. F.: *see* Kangasniemi, M.
Jaeger, D.: *see* Kotter, E.
Jentzen, W.: *see* Antoch, G.
Jin, G. Y.: *see* Lee, J. M.
Jorge-Costa, M.: *see* Montet, X.
Juergens, K. U.: *see* Grude, M.
Juha-Matti, I., Rami, H., Jukka, P., Osmo, T.: Accuracy of computer-aided measurements in endovascular stent-graft planning: An experimental study with two phantoms, 164
Jukka, P.: *see* Juha-Matti, I.
Jung, B.: *see* Mahnken, A. H.

K

Kachelriess, M.: *see* Achenbach, S.
Kainberger, F.: *see* Wagner, A.
Kaiser, W. A.: *see* Hansch, A.; Pfleiderer, S. O. R.
Kalender, W. A.: *see* Achenbach, S.
Kalinowski, M., Kress, O., Wels, T., Alfke, H., Klose, K. J., Wagner, H.-J.: 1-Molar gadobutrol as a contrast agent for computed tomog-

raphy: Results from a comparative porcine study, 193
Kanematsu, M.: *see* Murakami, T.
Kangasniemi, M., Stafford, R. J., Price, R. E., Jackson, E. F., Hazle, J. D.: Dynamic gadolinium uptake in thermally treated canine brain tissue and experimental cerebral tumors, 102
Kapell, S.: *see* Knollmann, F. D.
Kasper, S.: *see* Gruber, S.
Katoh, M.: *see* Wildberger, J. E.
Kauczor, H.-U.: *see* Gast, K. K.; Heussel, C. P.
Kavoussi, L. R.: *see* Solomon, S. B.
Kawakami, Y.: *see* Suga, K.
Kawata, S.: *see* Murakami, T.
Kemp, G. J.: *see* Gruber, S.
Kessler, W.: *see* Regenfus, M.
Kiessling, F., Krix, M., Heilmann, M., Vosseler, S., Lichy, M., Fink, C., Farhan, N., Kleinschmidt, K., Schad, L., Fusenig, N. E., Delorme, S.: Comparing dynamic parameters of tumor vascularization in nude mice revealed by magnetic resonance imaging and contrast-enhanced intermittent power doppler sonography, 516
Kim, C. J.: *see* Cheon, J.-E.
Kim, C. S.: *see* Kim, Y. K.; Lee, J. M.
Kim, H. J., Lee, C. H., Lee, S. H., Moon, T. Y.: Magnetic resonance imaging and histologic findings of experimental cerebral fat embolism, 625
Kim, H.-W., Buckley, D. L., Peterson, D. M., Duensing, G. R., Caserta, J., Fitzsimmons, J., Blackband, S. J.: In vivo prostate magnetic resonance imaging and magnetic resonance spectroscopy at 3 Tesla using a transceive pelvic phased array coil: Preliminary results, 443
Kim, I.-O.: *see* Cheon, J.-E.
Kim, K., Kim, S. H., Yang, C. W., Li, C., Chung, Y. A., Lee, S. Y., Sohn, H. S., Chung, S. K.: Differentiation between acute cyclosporine nephrotoxicity and acute tubular necrosis using enalaprilat renal scintigraphy in rats, 473
Kim, S. H.: *see* Kim, K.
Kim, S. J.: *see* Cha, S.-H.
Kim, S. W.: *see* Kim, Y. K.
Kim, T.: *see* Murakami, T.
Kim, T.-H.: *see* Shin, J. H.
Kim, W. S.: *see* Cheon, J.-E.
Kim, Y. H., Choi, B. I., Cho, W. H., Lim, S., Moon, W. K., Han, J. K., Weinmann, H.-J., Chang, K.-H.: Dynamic contrast-enhanced MR imaging of VX2 carcinomas after X-irradiation in rabbits: Comparison of gadopentetate dimeglumine and a macromolecular contrast agent, 539
Kim, Y. K., Lee, J. M., Kim, S. W., Kim, C. S.: Combined radiofrequency ablation and hot saline injection in rabbit liver, 725
Kirchin, M. A.: *see* Schneider, G.; Wikström, J.
Kivelitz, D., Wagner, S., Schnorr, J., Wetzler, R., Busch, M., Melzer, A., Taupitz, M., Hamm, B.: A vascular stent as an active component for locally enhanced magnetic resonance imaging: Initial in vivo imaging results after catheter-guided placement in rabbits, 147
Kleinschmidt, K.: *see* Kiessling, F.
Klose, K. J.: *see* Kalinowski, M.
Klutch, R.: *see* Sicotte, N. L.
Knez, A.: *see* Becker, C. R.
Knitz, F.: *see* Gast, K. K.
Knollmann, F. D., Helmig, K., Kapell, S., Hummel, M., Bocksch, W., Hetzer, R., Felix, R.: Coronary artery calcium scoring: Diagnostic accuracy of different software implementations, 761
Knopp, M. V.: *see* Wikström, J.
Kohno, M.: *see* Masugata, H.
Kölker, C.: *see* Spuentrup, E.
König, C. W., Trübenbach, J., Böhm, P., Fritz, J., Duda, S. H., Pereira, P. L.: Magnetic resonance-guided transcortical biopsy of bone marrow lesions using a magnetic resonance imaging-compatible piezoelectric power drill: Preliminary experience, 159
Kool, L. S.: *see* Schneider, G.
Kotter, E., Bley, T. A., Saueressig, U., Fisch, D., Springer, O., Winterer, J. T., Schaefer, O., Langer, M.: Comparison of the detectability of high- and low-contrast details on a TFT screen and a CRT screen designed for radiologic diagnosis, 719
Kotter, E., Roesner, A., Winterer, J. T., Ghanem, N., Einert, A., Jaeger, D., Uhrmeister, P., Langer, M.: Evaluation of lossy data compression of chest x-rays: A receiver operating characteristic study, 243
Krämer, S.: *see* Maintz, D.
Krause, W., Gilles, T.: A new animal model for testing renal tolerance of contrast agents, 704
Krause, W., Schromm, M.: Use of additives to contrast media to improve imaging in plain film and computed tomography urography, 17
Kreicbergs, A.: *see* Ahmad, T.
Kresnik, E.: *see* Gallowitsch, H.-J.
Kress, O.: *see* Kalinowski, M.
Krix, M.: *see* Kiessling, F.
Kröber, S.-M.: *see* Brieger, J.
Krombach, G. A.: *see* Spuentrup, E.
Kühl, H. P.: *see* Spuentrup, E.
Kumano, S.: *see* Murakami, T.
Kumnig, G.: *see* Gallowitsch, H.-J.
Kuo, T. Y., Skedros, J. G., Bloebaum, R. D.: Measurement of femoral anteversion by biplane radiography and computed tomography imaging: Comparison with an anatomic reference, 221
Kuzo, R. S.: *see* Gerber, T. C.
Kwan, O. L.: *see* Masugata, H.
Kwon, O-K., Han, M. H, Oh, C. W., Park, I. A., Choe, G., Lee, K. H., Chang, K.-H., Han, D. H.: Embolization with autologous fibroblast-attached platinum coils in canine carotid artery aneurysms: Histopathological differences from plain coil embolization, 281

L

Labarthe, B., Idee, J.-M., Burnett, R., Corot, C: In vivo comparative antithrombotic effects of ioxaglate and iohexol and interaction with the platelet antiaggregant clopidogrel, 34
Lafitte, S.: *see* Masugata, H.
Langenberger, H., Shimizu, Y., Windischberger, C., Grampp, S., Berg, A., Ferlitsch, K., Moser, E.: Bone homogeneity factor: An advanced tool for the assessment of osteoporotic bone structure in high-resolution magnetic resonance images, 467
Langer, M.: *see* Kotter, E.
Laub, G.: *see* Regenfus, M.
Lauenstein, T. C.: *see* Herborn, C. U.
Laukemper-Ostendorf, S.: *see* Heussel, C. P.
Laurent, S.: *see* Burtea, C.
Lauzon, M. L.: *see* Frayne, R.
Lawaczeck, R., Diekmann, F., Dieckmann, S., Hamm, B., Bick, U., Press, W.-R., Schirmer, H., Schön, K., Weinmann, H.-J.: New contrast media designed for x-ray energy subtraction imaging in digital mammography, 602
Leber, A.: *see* Becker, C. R.
Lee, C. H.: *see* Kim, H. J.
Lee, J.-H.: *see* Shin, J. H.
Lee, J. M.: *see* Kim, Y. K.
Lee, J. M., Jin, G. Y., Li, C. A., Chung, G. H., Lee, S. Y., Han, Y. M., Chung, M. J., Kim, C. S.: Percutaneous radiofrequency thermal ablation of lung VX2 tumors in a rabbit model using a cooled tip-electrode: Feasibility, safety, and effectiveness, 129
Lee, K. H.: *see* Kwon, O.-K.
Lee, S. H.: *see* Kim, H. J.
Lee, S. Y.: *see* Kim, K.; Lee, J. M.
Legras, P.: *see* Chapon, C.
Le Jeune, J.-J.: *see* Chapon, C.
Lemaire, L.: *see* Chapon, C.
Lenkinski, R. E.: *see* Alves, F. C.
Ley, S.: *see* Heussel, C. P.
Li, C.: *see* Kim, K.
Li, C. A.: *see* Lee, J. M.
Li, W.: *see* Storey, P.
Lichy, M.: *see* Kiessling, F.
Lichy, M. P.: *see* Weber, M.-A.
Lim, J-O.: *see* Shin, J. H.
Lim, S.: *see* Kim, Y. H.
Lind, P.: *see* Gallowitsch, H.-J.
Link, T. M.: *see* Boehm, H. F.
Liu, S. C. K.: *see* Mao, S.
Lodemann, K.-P.: *see* Schneider, G.
Lorenzen, J., Sinkus, R., Biesterfeldt, M., Adam, G.: Menstrual cycle dependence of breast parenchyma elasticity: Estimation with magnetic resonance elastography of breast tissue during the menstrual cycle, 236
Lubag, A. J. M.: *see* Alves, F. C.
Lubienski, A.: *see* Grenacher, L.
Luechinger, R.: *see* Spiegel, M. A.

M

Maas, R.: *see* Schneider, G.
Mahnken, A. H.: *see* Spuentrup, E.; Wildberger, J. E.
Mahnken, A. H., Raupach, R., Wildberger, J. E., Jung, B., Heussen, N., Flohr, T. G., Günther, R. W., Schaller, S.: A new algorithm for metal artifact reduction in computed tomography: In vitro and in vivo evaluation after total hip replacement, 769
Maintz, D., Seifarth, H., Flohr, T., Krämer, S., Wichter, T., Heindel, W., Fischbach, R.: Improved coronary artery stent visualization and in-stent stenosis detection using 16-slice computed-tomography and dedicated image reconstruction technique, 790
Majima, K.: *see* Tsuda, M.
Majumdar, S.: *see* Boehm, H. F.
Malich, A.: *see* Hansch, A.
Malich, A., Fuchsjäger, M.: Electrical impedance scanning in classifying suspicious breast lesions (letter), 302
Mao, S., Child, J., Carson, S., Liu, S. C. K., Oudiz, R. J., Budoff, M. J.: Sensitivity to detect small coronary artery calcium lesions with varying slice thickness using electron beam tomography, 183
Marchal, G.: *see* Wikström, J.
Marescaux, L.: *see* Chapon, C.
Markstaller, K.: *see* Gast, K. K.
Marx, C.: *see* Pfleiderer, S. O. R.
Masugata, H., DeMaria, A. N., Peters, B., Lafitte, S., Strachan, G. M., Kwan, O. L., Ohmori, K., Mizushige, K., Kohno, M.: Difference of optimal dose of contrast agent between gray-scale and power doppler imaging

in assessing graded coronary stenosis by myocardial contrast echocardiography, 550
Masugata, H., Peters, B., Lafitte, S., Yao, B., Strachan, M., Kwan, O. L., Ohmori, K., Mizushige, K., Kohno, M.: Comparison of open- and closed-chest canine model for quantification of coronary stenosis severity by myocardial contrast echocardiography, 44
Matsunaga, N.: *see* Suga, K.
Matula, C.: *see* Barth, M.
Max, M.: *see* Wildberger, J. E.
Mayer, E: *see* Gast, K. K.
Mazziotta, J. C.: *see* Sicotte, N. L.
McDonald, M.A., Watkin, K. L.: Small particulate gadolinium oxide and gadolinium oxide albumin microspheres as multimodal contrast and therapeutic agents, 305
Mehrain, S.: *see* Ba-Ssalamah, A.
Melzer, A.: *see* Kivelitz, D.
Merritt, M. E.: *see* Alves, F. C.
Meuwly, J.-Y., Thiran, J.-P., Gudinchet, F.: Application of adaptive image processing technique to real-time spatial compound ultrasound imaging improves image quality, 257
Mikosch, P.: *see* Gallowitsch, H.-J.
Miyachi, H.: *see* Tsuda, M.
Mizushige, K.: *see* Masugata, H.
Mlynárik, V.: *see* Barth, M.; Ba-Ssalamah, A.; Gruber, S.
Modai, D.: *see* Peer, A.
Monetti, R. A.: *see* Boehm, H. F.
Montet, X., Ivancevic, M. K., Belenger, J., Jorge-Costa, M., Pochon, S., Pechère, A., Terrier, F., Vallée, J.-P.: Noninvasive measurement of absolute renal perfusion by contrast medium-enhanced magnetic resonance imaging, 584
Moon, T. Y.: *see* Kim, H. J.
Moon, W. K.: *see* Kim, Y. H.
Morfill, G.: *see* Boehm, H. F.
Morin, R. L.: *see* Gerber, T. C.
Mortensen, J.: *see* Rawashdeh, Y. F.
Moser, E.: *see* Berg, A.; Gruber, S.; Langenberger, H.
Moser, E., Trattnig, S.: 3.0 Tesla MR systems, 375
Moshage, W.: *see* Regenfus, M.
Mueller, D.: *see* Boehm, H. F.
Mueller, S. P.: *see* Antoch, G.
Mukherji, S. K.: *see* Bui, C. D.
Muller, J. G.: *see* Grude, M.
Muller, R. N.: *see* Burtea, C.
Murakami, T., Kim, T., Kawata, S., Kanematsu, M., Federle, M. P., Hori, M., Okada, A., Kumano, S., Sugihara, E., Tomoda, K., Nakamura, H.: Evaluation of optimal timing of arterial phase imaging for the detection of hypervascular hepatocellular carcinoma by using triple arterial phase imaging with multidetector-row helical computed tomography, 497

N

Nakada, T.: Clinical experience on 3.0 T systems in Niigata, 1996 to 2002, 377
Nakamura, H.: *see* Murakami, T.
Naul, L. G.: *see* Desai, N.
Neves, M.: *see* Alves, F. C.
Newitt, D.: *see* Boehm, H. F.
Nguyen, T. H.: *see* Spuentrup, E.
Nicol, T. L.: *see* Solomon, S. B.
Nöbauer-Huhmann, I.-M.: *see* Barth, M.
Nöbauer-Huhmann, I. M.: *see* Ba-Ssalamah, A.
Nobuhiko, O.: *see* Suga, K.
Nolte-Ernsting, C., Glowinski, A., Schaeffter, T., Adam, G., Günther, R. W.: Gadolinium-enhanced magnetic resonance fluoroscopy used as micturating cystourethrography: Experiences in adult male patients, 617

O

O'Brien, P. C.: *see* Gerber, T. C.
Ogasawara, N.: *see* Suga, K.
Oh, C. W.: *see* Kwon, O.-K.
Ohlsson, C.: *see* Ahmad, T.
Ohmori, K.: *see* Masugata, H.
Oike, M.: *see* Sumimura, T.
Oishi, R.: *see* Sumimura, T.
Oka, M.: *see* Sumimura, T.
Okada, A.: *see* Murakami, T.
Okada, M.: *see* Suga, K.
Osmo, T.: *see* Juha-Matti, I.
Östenson, C.-G.: *see* Ahmad, T.
Ottoboni, T. B.: *see* Wisner, E. R.
Oudiz, R. J.: *see* Mao, S.
Oudkerk, M.: *see* Wikström, J.

P

Paetzel, C.: *see* Völk, M.
Papadokostakis, G.: *see* Damilakis, J.
Park, I. A.: *see* Kwon, O.-K.
Pastor, K.: *see* Gerber, T. C.
Patel, D.: *see* Wisner, E. R.
Pattynama, P. M. T.: *see* Wikström, J.
Paul, M.: *see* Grude, M.
Pechère, A.: *see* Montet, X.
Peer, A., Averbukh, Z., Berman, S., Modai, D., Averbukh, M., Weissgarten, J.: Contrast media augmented apoptosis of cultured renal mesangial, tubular, epithelial, endothelial, and hepatic cells, 177
Pereira, P. L.: *see* Brieger, J.; König, C. W.
Perisinakis, K.: *see* Damilakis, J.
Peters, B.: *see* Masugata, H.
Peterson, D. M.: *see* Kim, H.-W.
Peterson, D. M., Duensing, G. R., Caserta, J., Fitzsimmons, J. R.: An MR transceive phased array designed for spinal cord imaging at Tesla Preliminary Investigations of Spinal Cord Imaging at 3 T, 428
Pfleiderer, S. O. R., Reichenbach, J. R., Azhari, T., Marx, C., Wurdinger, S., Kaiser, W. A.: Dedicated double breast coil for magnetic resonance mammography imaging, biopsy, and preoperative localization, 1
Pinker, K.: *see* Ba-Ssalamah, A.
Pitts, J.: *see* Hinton, D. P.
Pochon, S.: *see* Montet, X.
Pohle, K.: *see* Achenbach, S.
Port, M.: *see* Corot, C.
Post, M.: *see* Storey, P.
Prasad, P. V.: *see* Storey, P.
Prata, M. I. M.: *see* Alves, F. C.
Press, W.-R.: *see* Lawaczeck, R.
Price, R. E.: *see* Kangasniemi, M.
Prokesch, R.: *see* Ba-Ssalamah, A.
Puderbach, M.: *see* Fink, C.
Puderbach, M. U.: *see* Gast, K. K.

Q

Quintel, M.: *see* Heussel, C. P.

R

Raeth, C.: *see* Boehm, H. F.
Rami, H.: *see* Juha-Matti, I.
Rand, T.: *see* Berg, A.
Raupach, R.: *see* Mahnken, A. H.
Rawashdeh, Y. F., Hørlyck, A., Mortensen, J., Hvistendahl, J. J., Frokiaer, J., Djurhuus, J. C.: Resistive index: An experimental study of acute complete unilateral ureteral obstruction, 153
Regenfus, M., Ropers, D., Achenbach, S., Schlundt, C., Kessler, W., Laub, G., Moshage, W., Daniel, W. G.: Diagnostic value of maximum intensity projections *versus* source images for assessment of contrast-enhanced three-dimensional breath-hold magnetic resonance coronary angiography, 200
Reiber, J. H. C.: *see* Box, F. M. A.
Reichenbach, J. R.: *see* Barth, M.; Pfleiderer, S. O. R.
Reiser, M. F.: *see* Becker, C. R.
Richter, G. M.: *see* Grenacher, L.
Rikimaru, H.: *see* Tsuda, M.
Robert, P.: *see* Corot, C.
Rodriguez, I.: *see* Gast, K. K.
Roesner, A.: *see* Kotter, E.
Ropers, D.: *see* Achenbach, S.; Regenfus, M.
Rossaint, R.: *see* Wildberger, J. E.
Ruehm, S. G.: *see* Herborn, C. U.
Rummeny, E.: *see* Boehm, H. F.; Schneider, G.
Runge, V. M.:, 802. *see* Desai, N.

S

Sailer, J.: *see* Berg, A.
Saint-André, J. P.: *see* Chapon, C.
Saito, H.: *see* Tsuda, M.
Salah, A.: *see* Chaabane, L.
Santos, A. C.: *see* Alves, F. C.
Saueressig, U.: *see* Kotter, E.
Sauner, D.: *see* Hansch, A.
Schad, L.: *see* Kiessling, F.
Schad, L. R.: *see* Weber, M.-A.
Schaefer, O.: *see* Kotter, E.
Schaeffter, T.: *see* Nolte-Ernsting, C.; Spuentrup, E.
Schaller, S.: *see* Mahnken, A. H.; Wildberger, J. E.
Scheller, B.: *see* Clauß, W.
Schenk, M.: *see* Brieger, J.
Schibany, N.: *see* Ba-Ssalamah, A.
Schicho, K.: *see* Wagner, A.
Schick, F.: *see* Brieger, J.
Schild, H.: *see* Böhm, I.
Schirmer, H.: *see* Lawaczeck, R.
Schlemmer, H.-P.: *see* Weber, M.-A.
Schlundt, C.: *see* Regenfus, M.
Schmähl, A.: *see* Fink, C.
Schmidt, D.: *see* Brieger, J.
Schmiedeskamp, J.: *see* Gast, K. K.
Schmitt, A.: *see* Clauß, W.
Schmitt, F.: *see* Hinton, D. P.
Schmittner, M.: *see* Heussel, C. P.
Schneider, G., Maas, R., Kool, L. S., Rummeny, E., Gehl, H.-B., Lodemann, K.-P., Kirchin, M. A.: Low-dose gadobenate dimeglumine *versus* standard dose gadopentetate dimeglumine for contrast-enhanced magnetic resonance imaging of the liver: An intra-individual crossover comparison, 85
Schnorr, J.: *see* Kivelitz, D.
Schoepf, U. J.: *see* Becker, C. R.
Schöggl, A.: *see* Barth, M.
Scholz, A.: *see* Heussel, C. P.
Schön, K.: *see* Lawaczeck, R.
Schreiber, W. G.: *see* Gast, K. K.; Heussel, C. P.
Schromm, M.: *see* Krause, W.
Schwitter, J.: *see* Spiegel, M. A.
Seifarth, H.: *see* Maintz, D.
Seitz, J.: *see* Völk, M.
Sendo, T.: *see* Sumimura, T.
Seo, T.-S.: *see* Shin, J. H.

Seoane, P. R.: *see* Storey, P.
Sevick, R. J.: *see* Frayne, R.
Shen, N.: *see* Wikström, J.
Sherry, A. D.: *see* Alves, F. C.
Shimizu, Y.: *see* Langenberger, H.
Shin, J. H., He, X., Lee, J.-H., Seo, T.-S., Lim, J.-O., Kim, T.-H., Song, H.-Y.: Newly designed multifunctional coil catheter for gastrointestinal intervention: Feasibility determined by experimental study in dogs, 796
Short, R. E.: *see* Wisner, E. R.
Shreve, P. D.: *see* Bui, C. D.
Sicotte, N. L., Voskuhl, R. R., Bouvier, S., Klutch, R., Cohen, M. S., Mazziotta, J. C.: Comparison of multiple sclerosis lesions at 1.5 and 3.0 Tesla, 423
Singer, T.: *see* Berg, A.
Sinkus, R.: *see* Lorenzen, J.
Skedros, J. G.: *see* Kuo, T. Y.
Sohn, H. S.: *see* Kim, K.
Solomon, S. B., Nicol, T. L., Chan, D. Y., Fjield, T., Fried, N., Kavoussi, L. R.: Histologic evolution of high-intensity focused ultrasound in rabbit muscle, 293
Song, H.-Y.: *see* Shin, J. H.
Soulas, E. C.: *see* Chaabane, L.
Sovak, M.: *see* Clauβ, W.
Speck, U.: *see* Böhm, I.; Clauβ, W.
Spiegel, M. A., Luechinger, R., Schwitter, J., Boesiger, P.: RingTag: Ring-shaped tagging for myocardial centerline assessment, 669
Spilt, A.: *see* Box, F. M. A.
Spinazzi, A.: *see* Wikström, J.
Springer, O.: *see* Kotter, E.
Spuentrup, E., Buecker, A., Stuber, M., Botnar, R., Nguyen, T. H., Börnert, P., Kölker, C., Günther, R. W.: Navigator-gated coronary magnetic resonance angiography using steady-state-free-precession: Comparison to standard T2-prepared gradient-echo and spiral imaging, 263
Spuentrup, E., Mahnken, A. H., Kühl, H. P., Krombach, G. A., Botnar, R. M., Wall, A., Schaeffter, T., Günther, R. W., Buecker, A.: Fast interactive real-time magnetic resonance imaging of cardiac masses using spiral gradient echo and radial steady-state free precession sequences, 288
Stadlbauer, A.: *see* Gruber, S.
Stafford, R. J.: *see* Kangasniemi, M.
Stattaus, J.: *see* Antoch, G.
Storey, P., Danias, P. G., Post, M., Li, W., Seoane, P. R., Harnish, P. P., Edelman, R. R., Prasad, P. V.: Preliminary evaluation of EVP 1001–1: A new cardiac-specific magnetic resonance contrast agent with kinetics suitable for steady-state imaging of the ischemic heart, 642
Strachan, G. M.: *see* Masugata, H.
Strachan, M.: *see* Masugata, H.
Strotzer, M.: *see* Hamer, O. W.; Völk, M.
Stuber, M.: *see* Spuentrup, E.
Stupar, V., Berthezène, Y., Canet, E., Tournier, H., Dupuich, D., Crémillieux, Y.: Helium3 polarization using spin exchange tehnique: Application to simultaneous pulmonary ventilation/perfusion imaging in small animals, 334
Suga, K., Ogasawara, N., Yuan, Y., Okada, M., Matsunaga, N., Tangoku, A.: Visualization of breast lymphatic pathways with an indirect computed tomography lymphography using a nonionic monometric contrast medium iopamidol: Preliminary results, 73
Suga, K., Yuan, Y., Nobuhiko, O., Okada, M., Kawakami, Y., Matsunaga, N.: Potential of magnetic resonance lymphography with intrapulmonary injection of gadopentetate dimeglumine for visualization of the pulmonary lymphatic basin in dogs: Preliminary results, 679
Suga, K., Yuan, Y., Ogasawara, N., Okada, M., Matsunaga, N.: Visualization of normal and interrupted lymphatic drainage in dog legs with interstitial MR lymphography using an extracellular MR contrast agent, gadopentetate dimeglumine, 349
Sugihara, E.: *see* Murakami, T.
Sumimura, T., Sendo, T., Itoh, Y., Oka, M., Oike, M., Ito, Y., Oishi, R.: Calcium-dependent injury of human microvascular endothelial cells induced by a variety of iodinated radiographic contrast media, 366

T

Takahashi, S.: *see* Tsuda, M.
Tangoku, A.: *see* Suga, K.
Taupitz, M.: *see* Kivelitz, D.
Terrier, F.: *see* Montet, X.
Thelen, M.: *see* Gast, K. K.; Heussel, C. P.
Thilmann, C.: *see* Weber, M.-A.
Thiran, J.-P.: *see* Meuwly, J.-Y.
Tomoda, K.: *see* Murakami, T.
Tournier, H.: *see* Stupar, V.
Trattnig, S.: *see* Barth, M.; Ba-Ssalamah, A.; Moser, E.
Trübenbach, J.: *see* Brieger, J.; König, C. W.
Tsagaraki, I.: *see* Damilakis, J.
Tsuda, M., Rikimaru, H., Majima, K., Yamada, T., Saito, H., Ishibashi, T., Takahashi, S., Miyachi, H., Endoh, M., Yamada, S.: Time-related changes of radiofrequency ablation lesion in the normal rabbit liver: Findings of magnetic resonance imaging and histopathology, 525

U

Uhrmeister, P.: *see* Kotter, E.
Ulzheimer, S.: *see* Achenbach, S.

V

Vallée, J.-P.: *see* Montet, X.
Van Buchem, M. A.: *see* Box, F. M. A.
Vander Elst, L.: *see* Burtea, C.
van der Geest, R. J.: *see* Box, F. M. A.
Violas, X.: *see* Corot, C.
Völk, M.: *see* Hamer, O. W.
Völk, M., Angele, P., Hamer, O., Feuerbach, S., Strotzer, M.: Digital image composition in long-leg radiography with a flat-panel detector: First clinical experiences, 189
Völk, M., Paetzel, C., Angele, P., Seitz, J., Füchtmeier, B., Hente, R., Feuerbach, S., Strotzer, M.: Routine skeleton radiography using a flat-panel detector: Image quality and clinical acceptance at 50% dose reduction, 230
Voskuhl, R. R.: *see* Sicotte, N. L.
Vosseler, S.: *see* Kiessling, F.
Vrahoriti, H.: *see* Damilakis, J.

W

Wagner, A., Schicho, K., Kainberger, F., Birkfellner, W., Grampp, S., Ewers, R.: Quantification and clinical relevance of head motion during computed tomography, 733
Wagner, H.-J.: *see* Kalinowski, M.
Wagner, S.: *see* Kivelitz, D.
Wald, L. L.: *see* Hinton, D. P.
Wall, A.: *see* Spuentrup, E.
Wang, S. J., Brechbiel, M., Wiener, E. C.: Characteristics of a new MRI contrast agent prepared from polypropyleneimine dendrimers, generation 2, 662
Wasser, M. N.: *see* Wikström, J.
Watkin, K. L.: *see* McDonald, M. A.
Weber, M.-A., Günther, M., Lichy, M. P., Delorme, S., Bongers, A., Thilmann, C., Essig, M., Zuna, I., Schad, L. R., Debus, J., Schlemmer, H.-P.: Comparison of arterial spin-labeling techniques and dynamic susceptibility-weighted contrast-enhanced MRI in perfusion imaging of normal brain tissue, 712
Weiler, N.: *see* Gast, K. K.; Heussel, C. P.
Wein, B. B.: *see* Wildberger, J. E.
Weinmann, H.-J.: *see* Kim, Y. H.; Lawaczeck, R.
Weiss, C.: *see* Wildberger, J. E.
Weissgarten, J.: *see* Peer, A.
Wels, T.: *see* Kalinowski, M.
Wenkel, E.: *see* Achenbach, S.
Wersebe, A.: Electrical impedance scanning in classifying suspicious breast lesions (reply to letter), 303
Wetzler, R.: *see* Kivelitz, D.
Wichter, T.: *see* Grude, M.; Maintz, D.
Wiener, E. C.: *see* Wang, S. J.
Wikström, J., Wasser, M. N., Pattynama, P. M. T., Bonomo, L., Hamm, B., Del Maschio, A., Knopp, M. V., Marchal, G., Barentsz, J. O., Oudkerk, M., Hentrich, H.-R., Daprà, M., Kirchin, M. A., Shen, N., Spinazzi, A., Ahlström, H.: Gadobenate dimeglumine-enhanced magnetic resonance angiography of the pelvic arteries, 504
Wildberger, J. E.: *see* Mahnken, A. H.
Wildberger, J. E., Max, M., Wein, B. B., Mahnken, A. H., Weiss, C., Dembinski, R., Katoh, M., Schaller, S., Rossaint, R., Günther, R. W.: Low-dose multislice spiral computed tomography in acute lung injury: Animal experience, 9
Windischberger, C.: *see* Langenberger, H.
Winterer, J. T.: *see* Kotter, E.
Wisner, E. R., Ferrara, K. W., Short, R. E., Ottoboni, T. B., Gabe, J. D., Patel, D.: Sentinel node detection using contrast-enhanced power doppler ultrasound lymphography, 358
Wurdinger, S.: *see* Pfleiderer, S. O. R.

Y

Yamada, S.: *see* Tsuda, M.
Yamada, T.: *see* Tsuda, M.
Yang, C. W.: *see* Kim, K.
Yao, B.: *see* Masugata, H.
Yeon, K. M.: *see* Cheon, J.-E.
Yoo, W. J.: *see* Cheon, J.-E.
Yoon, C. J.: *see* Cha, S.-H.
Yuan, Y.: *see* Suga, K.

Z

Zaheer, A.: *see* Alves, F. C.
Zhang, S.: *see* Alves, F. C.
Zorger, N.: *see* Hamer, O. W.
Zuna, I.: *see* Weber, M.-A.

Subject Index—Volume 38

A

Adaptive image processing technique (XRES), applied to US images, 257
Allergy, cytokine profiles after nonionic dimeric contrast medium injection, 776
Aneurysms, embolization with autologous fibroblast-attached platinum coils in canine carotid artery aneurysms, 281
Angiogenesis, tumor vascularization studies with dynamic MRI and Doppler US (mouse model), 516
Angiography, magnetic resonance
  comparison of Gd-DTPA, Gd-BOPTA, and galobutrol for pelvic 3D MRA, 27
  coronary, maximum intensity projections for assessment of, 200
  different types of blood pool agents in rabbit MRA-like protocol, 311
  Gd-BOPTA–enhanced MRA of pelvic arteries, 504
  navigator-gated coronary MRA using steady-state-free-precession, 263
  new glucosylated derivatives of Gd-DTPA for, 320
Animal models. *See* specific species
Aorta
  accuracy of computer-aided measurements in stent-graft planning, 164
  high-resolution MRI of atherosclerotic lesions at 2 Tesla (mouse model), 532
  resistance to hoop stress in balloon expandable stents, 65
  vascular stent as active component for locally enhanced MRI (rabbit model), 147
Arthritis
  diagnosis by autofluorescence (mouse model), 578
  diffusivity imaging at 3 T for detection of degenerative changes, 460
Atherosclerosis
  detecting small coronary artery calcium lesions using electron beam tomography, 183
  diffusivity and T_2 imaging at 3 T for detection of degenerative changes in atherosclerotic arteries, 452
  high-resolution MRI of atherosclerotic lesions at 2 Tesla (mouse model), 532
  reconstruction windows for multislice CT to quantify coronary calcium, 108 (erratum, 241)
  resistance to hoop stress in balloon expandable stents, 65
Autofluorescence, diagnosis of arthritis by (mouse model), 578

B

Biopsy
  double breast coil for MR mammography, 1
  MR-guided biopsy of bone marrow lesions using piezoelectric power drill, 159
Blood flow. *See also* Perfusion
  measurement of absolute renal perfusion by MRI (rabbit model), 584
  quantification of blood flow in small vessels with velocity encoded MRI, 567
Blood-pool agents
  comparison of different types in rabbit MRA-like protocol, 311
  new glucosylated derivatives of Gd-DTPA for MRA, 320
Bone
  bone homogeneity factor for osteoporosis in high-resolution MRI, 467
  digital image composition in long-leg radiography with flat-panel detector, 189
  femoral anteversion measured by biplane radiography vs. CT, 221
  imaging findings after fat graft interposition in injured growth plate (rabbit model), 695
  MR-guided biopsy of bone marrow lesions using piezoelectric power drill, 159
  multisite quantitative US device in normal and osteoporotic postmenopausal women, 207
  peripheral quantitative CT for detection of diabetic osteopathy (rat model), 171
  skeletal radiography using flat-panel detector, image quality and clinical acceptance at 50% dose reduction, 230
  3D scaling method for predicting bone strength from high-resolution MRI, 269
Brain
  arterial spin-labeling techniques vs. dynamic susceptibility-weighted contrast-enhanced MRI in perfusion imaging of normal brain tissue, 712
  blood oxygenation level-dependent MRV of brain tumors at 3 Tesla, 409
  challenges and advantages of MRI at 3.0 Tesla, 385
  dynamic gadolinium uptake in thermally treated canine brain, 102
  effect of contrast dose and field strength in MR detection of brain metastases, 415
  MRI and histologic findings of cerebral fat embolism (cat model), 625
  MRI in glutaric acidemia type I, 489
  quantification of metabolic differences in frontal brain in depression by ^{1}H-MRS, 403
Breast
  double breast coil for MR mammography, 1
  electrical impedance scanning to classify suspicious lesions (letter and reply), 302
  F-18 FDG PET in diagnosis of breast tumor recurrence and metastases, 250
  menstrual-cycle related breast parenchyma elasticity, 236
  new contrast media for x-ray subtraction imaging in digital mammography, 602
  visualization of breast lymphatics with indirect CT lymphography using iopamidol, 73

C

Calcium
  calcium-dependent endothelial cell injury induced by contrast media, 366
  calibration phantoms and scoring thresholds for quantification of coronary calcium, 559
  detecting small coronary artery calcium lesions using electron beam tomography, 183
  diagnostic accuracy of software for coronary artery calcium scoring, 761
  reconstruction windows for multislice CT to quantify coronary calcium, 108 (erratum, 241)
Carotid arteries
  embolization with autologous fibroblast-attached platinum coils in canine carotid artery aneurysms, 281
  vascular responses to various self-expanding stents (dog model), 95
Cartilage, diffusivity imaging for detection of degenerative changes in arthritis, 460
Cat, MRI and histologic findings of cerebral fat embolism, 625
Catheters, multifunctional coil catheter for gastrointestinal intervention (dog model), 796
Cerebral fat embolism, MRI and histologic findings (cat model), 625
Chest radiography
  lossy data compression on, 243
  monitor luminescence and observer performance for detection of abnormalities, 57
Clopidogrel, antithrombotic effects of ioxaglate and iohexol and interaction with (rat model), 34

Coils
double breast coil for MR mammography, 1
embolization with autologous fibroblast-attached platinum coils in canine carotid artery aneurysms, 281
multifunctional coil catheter for gastrointestinal intervention (dog model), 796
prostate MRI and MRS using transceive pelvic phased array coil, 443
Comparative studies
amorphous silicon, flat-panel, x-ray detector vs. storage phosphor-based computed radiography, 212
antithrombotic effects of ioxaglate and iohexol and interaction with clopidogrel (rat model), 34
arterial spin-labeling techniques vs. dynamic susceptibility-weighted contrast-enhanced MRI in perfusion imaging of normal brain tissue, 712
comparison of Gd-DTPA, Gd-BOPTA, and galobutrol for pelvic 3D MRA, 27
coronary-artery visualization by electron beam tomography vs. multislice spiral CT, 119
detectability of high- and low-contrast details on a TFT screen and a CRT screen designed for radiologic diagnosis, 719
diagnostic accuracy of software for coronary artery calcium scoring, 761
different types of blood pool agents in rabbit MRA-like protocol, 311
femoral anteversion measured by biplane radiography vs. CT, 221
low-dose Gd-BOPTA vs. standard dose Gd-DTPA for liver MRI, 85
MRI vs. US to detect posterior tibialis tendon tears, 51
multiple sclerosis lesions at 1.5 and 3 Tesla, 423
1-molar gabobutrol as contrast agent for CT (pig model), 193
Computed tomography (CT)
accuracy of computer-aided measurements in stent-graft planning, 164
algorithm for metal artifact reduction: evaluation after total hip replacement, 769
calibration phantoms and scoring thresholds for quantification of coronary calcium, 559
coronary artery stent visualization and in-stent stenosis detection using 16-slice CT, 790
coronary-artery visualization by electron beam tomography vs. multislice spiral CT, 119
diagnostic accuracy of software for coronary artery calcium scoring, 761
effect of oral contrast agents in dual-modality PET/CT imaging, 784
electron beam (*See* Electron beam tomography)
femoral anteversion measured by biplane radiography vs. CT, 221
global left ventricular function evaluated with multidetector CT compared with MRI, 653
low-dose multislice spiral CT in acute lung injury (pig model), 9
1-molar gabobutrol as contrast agent for (pig model), 193
optimal contrast application for cardiac 4-detector-row CT, 690
optimal timing of helical CT arterial phase imaging in hypervascular hepatocellular carcinoma, 497
percutaneous radiofrequency thermal ablation of lung tumors (rabbit model), 129
peripheral quantitative CT for detection of diabetic osteopathy (rat model), 171
quantification and clinical relevance of head motion during, 733
reconstruction windows for multislice CT to quantify coronary calcium, 108 (erratum, 241)
small particulate gadolinium oxide and gadolinium oxide albumin microspheres as multimodal contrast agents, 305
use of additives to contrast media to improve imaging in plain film and CT urography (rabbit and dog models), 17
visualization of breast lymphatics with indirect CT lymphography using iopamidol, 73
Computer applications
accuracy of computer-aided measurements in stent-graft planning, 164
detectability of high- and low-contrast details on a TFT screen and a CRT screen designed for radiologic diagnosis, 719
diagnostic accuracy of software for coronary artery calcium scoring, 761
lossy data compression on chest radiography, 243
monitor luminescence and observer performance for detection of abnormalities on chest radiographs, 57
Contrast agents
antithrombotic effects of ioxaglate and iohexol and interaction with clopidogrel (rat model), 34
arterial spin-labeling techniques vs. dynamic susceptibility-weighted contrast-enhanced MRI in perfusion imaging of normal brain tissue, 712
calcium-dependent endothelial cell injury induced by, 366
comparative study of 1-molar gabobutrol for CT (pig model), 193
comparison of Gd-DTPA, Gd-BOPTA, and galobutrol for pelvic 3D MRA, 27
contrast media augmented apoptosis of cultured renal and hepatic cells, 177
cytokine profiles after nonionic dimeric contrast medium injection, 776
dose in assessing coronary stenosis by myocardial contrast echocardiography (dog model), 550
dynamic contrast-enhanced MRI of VX2 carcinomas in rabbits, 539
effect of contrast dose and field strength in MR detection of brain metastases, 415
effect of oral contrast agents in dual-modality PET/CT imaging, 784
effect on neointimal proliferation and restenosis after coronary stenting in pigs, 743
Gd-BOPTA–enhanced MRA of pelvic arteries, 504
low-dose Gd-BOPTA vs. standard dose Gd-DTPA for liver MRI, 85
magnetic resonance research (editorial), 802
new animal model for testing renal tolerance of (dog model), 704
new glucosylated derivatives of Gd-DTPA for MRA, 320
new media for x-ray subtraction imaging in digital mammography, 602
new MRI contrast agent prepared from polypropyleneimine dendrimers, 662
optimal contrast application for cardiac 4-detector-row CT, 690
preliminary evaluation of EVP 1001–1, 642
sentinel node detection with contrast-enhanced Doppler US lymphography (dog model), 358
silencing of phosphonate-gadolinium MRI contrast by hydroxyapatite binding (rabbit model), 750
small particulate gadolinium oxide and gadolinium oxide albumin microspheres as multimodal agents, 305
use of additives to improve imaging in plain film and CT urography (rabbit and dog models), 17
visualization of breast lymphatics with indirect CT lymphography using iopamidol, 73
visualization of lymphatic drainage in dog legs with interstitial MR lymphography using Gd-DTPA, 349
Coronary arteries
calibration phantoms and scoring thresholds for quantification of coronary calcium, 559
coronary artery stent visualization and in-stent stenosis detection using 16-slice CT, 790
coronary-artery visualization by electron beam tomography vs. multislice spiral CT, 119
detecting small coronary artery calcium lesions using electron beam tomography, 183

diagnostic accuracy of software for coronary artery calcium scoring, 761
dose of contrast agent in assessing coronary stenosis by myocardial contrast echocardiography (dog model), 550
effect on neointimal proliferation and restenosis after coronary stenting in pigs, 743
maximum intensity projections for assessment of MR angiography, 200
navigator-gated coronary MRA using steady-state-free-precession, 263
open- vs. closed-chest canine model for quantification of coronary stenosis severity, 44
reconstruction windows for multislice CT to quantify coronary calcium, 108 (erratum, 241)
Culture studies
calcium-dependent endothelial cell injury induced by contrast media, 366
contrast media augmented apoptosis of cultured renal and hepatic cells, 177
Cyclosporine, enalaprilat renal scintigraphy to differentiate acute tubular necrosis from nephrotoxicity of, 473
Cystourethrography, micturating, gadolinium-enhanced MR fluoroscopy used as, 617
Cytokine profiles after nonionic dimeric contrast medium injection, 776

D

Depression, quantification of metabolic differences in frontal brain by ^{1}H-MRS in, 403
Diabetic osteopathy, peripheral quantitative CT for detection (rat model), 171
Dog
carotid artery responses to various self-expanding stents, 95
dose of contrast agent in assessing coronary stenosis by myocardial contrast echocardiography, 550
dynamic gadolinium uptake in thermally treated brain, 102
embolization with autologous fibroblast-attached platinum coils in canine carotid artery aneurysms, 281
multifunctional coil catheter for gastrointestinal intervention, 796
new animal model for testing renal tolerance of contrast agents, 704
open- vs. closed-chest canine model for quantification of coronary stenosis severity, 44
potential of MR lymphography with intrapulmonary injection of Gd-DTPA for visualization of pulmonary lymphatic basin, 679
sentinel node detection with Doppler US lymphography, 358
use of additives to improve imaging in plain film and CT urography, 17
visualization of lymphatic drainage in dog legs with interstitial MR lymphography using Gd-DTPA, 349
Dual energy x-ray absorptiometry (DEXA)
multisite quantitative US device in normal and osteoporotic postmenopausal women, 207
peripheral quantitative CT for detection of diabetic osteopathy (rat model), 171

E

Echocardiography, myocardial contrast
dose of contrast agent in assessing coronary stenosis by (dog model), 550
open- vs. closed-chest canine model for quantification of coronary stenosis severity, 44
Electrical impedance scanning, to classify suspicious breast lesions (letter and reply), 302
Electron beam tomography (EBT)
detection of small coronary artery calcium lesions, 183
vs. multislice spiral CT for coronary-artery visualization, 119
Embolization
with autologous fibroblast-attached platinum coils in canine carotid artery aneurysms, 281
MRI and histologic findings of cerebral fat embolism (cat model), 625
Endothelial cells, calcium-dependent injury induced by contrast media, 366
Errata, 241

F

Fat embolism, cerebral, MRI and histologic findings (cat model), 625
Fat graft interposition in injured growth plate (rabbit model), 695
Flat-panel detector
amorphous silicon, flat-panel, x-ray detector vs. storage phosphor-based computed radiography, 212
digital image composition in long-leg radiography with, 189
skeletal radiography image quality and clinical acceptance at 50% dose reduction, 230
Fluoroscopy, gadolinium-enhanced MR fluoroscopy used as micturating cystourethrography, 617

G

Gadobenate dimeglumine (Gd-BOPTA)
compared with Gd-DTPA and gadobutrol for pelvic 3D MRA, 27
Gd-BOPTA–enhanced MRA of pelvic arteries, 504
low-dose Gd-BOPTA vs. standard dose Gd-DTPA for liver MRI, 85
Gadobutrol
compared with Gd-DTPA and Gd-BOPTA for pelvic 3D MRA, 27
1-molar gabobutrol as contrast agent for (pig model), 193
Gadolinium diethylenetriaminepentaacetic acid (Gd-DTPA), new glucosylated derivatives of Gd-DTPA for MRA, 320
Gadolinium oxide, small particulate gadolinium oxide and gadolinium oxide albumin microspheres as multimodal contrast agents, 305
Gadomer-17, dynamic contrast-enhanced MRI of VX2 carcinomas in rabbits, 539
Gadopentetate dimeglumine (Gd-DTPA)
compared with Gd-BOPTA and gadobutrol for pelvic 3D MRA, 27
dynamic contrast-enhanced MRI of VX2 carcinomas in rabbits, 539
dynamic gadolinium uptake in thermally treated canine brain, 102
low-dose Gd-BOPTA vs. standard dose Gd-DTPA for liver MRI, 85
potential of MR lymphography with intrapulmonary injection of Gd-DTPA for visualization of pulmonary lymphatic basin in dogs, 679
for visualization of lymphatic drainage in dog legs with interstitial MR lymphography, 349
Gastrointestinal intervention, multifunctional coil catheter for (dog model), 796
Glutaric acidemia type I, MRI of brain in, 489
Growth plate injury, imaging findings after fat graft interposition in (rabbit model), 695

H

Head and neck tumors, nonsquamous, diagnostic accuracy of FDG-PET, 593
Head motion during CT, quantification and clinical relevance, 733
Heart
cardiac MRI on 1.5 and 3 T clinical whole body systems, 436
fast interactive real-time MRI of cardiac masses, 288
global left ventricular function evaluated with multidetector CT compared with MRI, 653

MRI evaluation of superparamagnetic iron oxide nanoparticles in rat myocardial infarction, 141
optimal contrast application for cardiac 4-detector-row CT, 690
preliminary evaluation of EVP 1001–1, 642
RingTag for myocardial centerline assessment, 669
Helium3
distribution of ventilation in lung transplant recipients evaluated by helium3 MRI, 341
spin-exchange polarization applied to ventilation/perfusion imaging (rat model), 334
Hepatocellular carcinoma, hypervascular, optimal timing of helical CT arterial phase imaging, 497
Histamine, cytokine profiles after nonionic dimeric contrast medium injection, 776
Histology
of high-intensity focused US in rabbit muscle, 293
MRI and histologic findings of cerebral fat embolism (cat model), 625
time-related MRI and histopathological changes in radiofrequency ablation lesion in rabbit liver, 525

I

Image quality
algorithm for metal artifact reduction in CT: evaluation after total hip replacement, 769
amorphous silicon, flat-panel, x-ray detector vs. storage phosphor-based computed radiography, 212
application of adaptive image processing technique to US images, 257
comparison of Gd-DTPA, Gd-BOPTA, and galobutrol for pelvic 3D MRA, 27
coronary-artery visualization by electron beam tomography vs. multislice spiral CT, 119
detectability of high- and low-contrast details on a TFT screen and a CRT screen designed for radiologic diagnosis, 719
digital image composition in long-leg radiography with flat-panel detector, 189
lossy data compression on chest radiography, 243
skeletal radiography using flat-panel detector, image quality and clinical acceptance at 50% dose reduction, 230
use of additives to improve imaging in plain film and CT urography (rabbit and dog models), 17
Interleukins, cytokine profiles after nonionic dimeric contrast medium injection, 776
Interventional procedures
combined radiofrequency ablation and hot saline injection in rabbit liver, 725
embolization with autologous fibroblast-attached platinum coils in canine carotid artery aneurysms, 281
in vivo efficiency of four radiofrequency ablation systems in pig liver, 609
multifunctional coil catheter for gastrointestinal intervention (dog model), 796
percutaneous radiofrequency thermal ablation of lung tumors (rabbit model), 129
Iohexol, antithrombotic effects compared with ioxaglate and interaction with clopidogrel (rat model), 34
Iopamidol, visualization of breast lymphatics with indirect CT lymphography using, 73
Ioxaglate, antithrombotic effects compared with iohexol and interaction with clopidogrel (rat model), 34
Iron oxide, MRI evaluation of superparamagnetic iron oxide nanoparticles in rat myocardial infarction, 141

K

Kidney
contrast media augmented apoptosis of cultured renal and hepatic cells, 177
enalaprilat renal scintigraphy to differentiate cyclosporine toxicity from acute tubular necrosis, 473
measurement of absolute renal perfusion by MRI (rabbit model), 584
new animal model for testing renal tolerance of contrast agents (dog model), 704
resistive index in complete unilateral ureteral obstruction (pig model), 153

L

Liver
combined radiofrequency ablation and hot saline injection in rabbit liver, 725
contrast media augmented apoptosis of cultured renal and hepatic cells, 177
in vivo efficiency of four radiofrequency ablation systems in pig liver, 609
low-dose Gd-BOPTA vs. standard dose Gd-DTPA for liver MRI, 85
optimal timing of helical CT arterial phase imaging in hypervascular hepatocellular carcinoma, 497
time-related MRI and histopathological changes in radiofrequency ablation lesion in rabbit liver, 525
Lung
distribution of ventilation in lung transplant recipients evaluated by helium3 MRI, 341
helium3 spin-exchange polarization applied to ventilation/perfusion imaging (rat model), 334
low-dose multislice spiral CT in acute lung injury (pig model), 9
^{19}F-MRI measurement of alveolar pO_2 in partial liquid ventilation, 635
partially parallel 3D MRI to assess lung perfusion, 482
percutaneous radiofrequency thermal ablation of lung tumors (rabbit model), 129
Lymphography
Doppler US, for sentinel node detection (dog model), 358
potential of MR lymphography with intrapulmonary injection of Gd-DTPA for visualization of pulmonary lymphatic basin in dogs, 679
visualization of breast lymphatics with indirect CT lymphography using iopamidol, 73
visualization of lymphatic drainage in dog legs with interstitial MR lymphography using Gd-DTPA, 349

M

Magnetic resonance angiography (MRA)
comparison of Gd-DTPA, Gd-BOPTA, and galobutrol for pelvic 3D MRA, 27
coronary, maximum intensity projections for assessment of, 200
different types of blood pool agents in rabbit MRA-like protocol, 311
Gd-BOPTA–enhanced MRA of pelvic arteries, 504
navigator-gated coronary MRA using steady-state-free-precession, 263
new glucosylated derivatives of Gd-DTPA for, 320
Magnetic resonance imaging (MRI)
arterial spin-labeling techniques vs. dynamic susceptibility-weighted contrast-enhanced MRI in perfusion imaging of normal brain tissue, 712
of brain in glutaric acidemia type I, 489
distribution of ventilation in lung transplant recipients evaluated by helium3 MRI, 341
double breast coil for MR mammography, 1

dynamic contrast-enhanced MRI of VX2 carcinomas in rabbits, 539
dynamic gadolinium uptake in thermally treated canine brain, 102
evaluation of superparamagnetic iron oxide nanoparticles in rat myocardial infarction, 141
fast interactive real-time MRI of cardiac masses, 288
gadolinium-enhanced MR fluoroscopy used as micturating cystourethrography, 617
global left ventricular function evaluated with multidetector CT compared with MRI, 653
high-resolution MRI of atherosclerotic lesions at 2 Tesla (mouse model), 532
imaging findings after fat graft interposition in injured growth plate (rabbit model), 695
low-dose Gd-BOPTA vs. standard dose Gd-DTPA for liver MRI, 85
magnetic resonance research (editorial), 802
for measurement of absolute renal perfusion (rabbit model), 584
menstrual-cycle related breast parenchyma elasticity, 236
MR-guided biopsy of bone marrow lesions using piezoelectric power drill, 159
MRI and histologic findings of cerebral fat embolism (cat model), 625
^{19}F-MRI measurement of alveolar pO_2 in partial liquid ventilation, 635
new contrast agent prepared from polypropyleneimine dendrimers, 662
partially parallel 3D MRI to assess lung perfusion, 482
potential of MR lymphography with intrapulmonary injection of Gd-DTPA for visualization of pulmonary lymphatic basin in dogs, 679
preliminary evaluation of EVP 1001–1, 642
quantification of blood flow in small vessels with velocity encoded MRI, 567
RingTag for myocardial centerline assessment, 669
silencing of phosphonate-gadolinium MRI contrast by hydroxyapatite binding (rabbit model), 750
small particulate gadolinium oxide and gadolinium oxide albumin microspheres as multimodal contrast agents, 305
at 3.0 Tesla
blood oxygenation level-dependent MRV of brain tumors, 409
bone homogeneity factor in osteoporosis, 467
cardiac MRI on 1.5 and 3 T clinical whole body systems, 436
challenges and advantages in neurological imaging, 385
clinical experience in Niigata, 1996 to 2002, 377
compared with multiple sclerosis lesions at 1.5 Tesla, 423
diffusivity and T_2 imaging for detection of degenerative changes in atherosclerotic arteries, 452
diffusivity imaging for detection of degenerative changes in arthritis, 460
editorial, 375
effect of contrast dose and field strength in detection of brain metastases, 415
MR transceive phased array designed for spinal cord imaging, 428
multiple sclerosis lesions at 1.5 and 3 Tesla, 423
prostate MRI and MRS using transceive pelvic phased array coil, 443
quantification of metabolic differences in frontal brain in depression by ^{1}H-MRS, 403
3D scaling method for predicting bone strength from high-resolution MRI, 269
time-related MRI and histopathological changes in radiofrequency ablation lesion in rabbit liver, 525
tumor vascularization studies with dynamic MRI and Doppler US (mouse model), 516
vs. US to detect posterior tibialis tendon tears, 51
visualization of lymphatic drainage in dog legs with interstitial MR lymphography using Gd-DTPA, 349
Magnetic resonance spectroscopy (MRS)
prostate MRI and MRS at 3 Tesla using transceive pelvic phased array coil, 443
quantification of metabolic differences in frontal brain in depression by ^{1}H-MRS, 403
Magnetic resonance venography (MRV), blood oxygenation level-dependent MRV of brain tumors at 3 Tesla, 409
Mammography. *See also* Breast
digital, new contrast media for x-ray subtraction imaging in, 602
MR, double breast coil for, 1
Menstrual-cycle related breast parenchyma elasticity, 236
Metal artifact reduction in CT, algorithm for, 769
Metastatic disease
diagnostic accuracy of FDG-PET in nonsquamous tumors of head and neck, 593
effect of contrast dose and field strength in MR detection of brain metastases, 415
F-18 FDG PET in diagnosis of breast tumor recurrence and metastases, 250
sentinel node detection with Doppler US lymphography (dog model), 358
Micturating cystourethrography, gadolinium-enhanced MR fluoroscopy used as, 617
Mouse
diagnosis of arthritis by autofluorescence, 578
high-resolution MRI of atherosclerotic lesions at 2 Tesla, 532
tumor vascularization studies with dynamic MRI and Doppler US, 516
MS325, different types of blood pool agents in rabbit MRA-like protocol, 311
Multiple sclerosis, comparison of lesions at 1.5 and 3 Tesla MRI, 423
Muscle, histologic evolution of high-intensity focused US in rabbit muscle, 293
Myocardial contrast echocardiography
dose of contrast agent in assessing coronary stenosis by (dog model), 550
open- vs. closed-chest canine model for quantification of coronary stenosis severity, 44
Myocardial infarction, MRI evaluation of superparamagnetic iron oxide nanoparticles in (rat model), 141

N

Neointimal hyperplasia
carotid artery responses to various self-expanding stents (dog model), 95
effect on neointimal proliferation and restenosis after coronary stenting in pigs, 743

O

Osteoporosis
bone homogeneity factor in high-resolution MRI, 467
multisite quantitative US device in normal and osteoporotic postmenopausal women, 207
Oxygen, ^{19}F-MRI measurement of alveolar pO_2 in partial liquid ventilation, 635

P

P792, different types of blood pool agents in rabbit MRA-like protocol, 311
Pelvis

comparison of Gd-DTPA, Gd-BOPTA, and galobutrol for pelvic 3D MRA, 27
Gd-BOPTA–enhanced MRA of pelvic arteries, 504
Perfusion. *See also* Blood flow
arterial spin-labeling techniques vs. dynamic susceptibility-weighted contrast-enhanced MRI in perfusion imaging of normal brain tissue, 712
helium3 spin-exchange polarization applied to ventilation/perfusion imaging (rat model), 334
measurement of absolute renal perfusion by MRI (rabbit model), 584
partially parallel 3D MRI to assess lung perfusion, 482
Phosphonate-gadolinium, silencing by hydroxyapatite binding (rabbit model), 750
Piezoelectric power drill, for MR-guided biopsy of bone marrow lesions, 159
Pig
effect of contrast media on neointimal proliferation and restenosis after coronary stenting, 743
in vivo efficiency of four radiofrequency ablation systems in pig liver, 609
low-dose multislice spiral CT in acute lung injury, 9
1-molar gabobutrol as contrast agent for CT, 193
Positron emission tomography (PET)
diagnostic accuracy of FDG-PET in nonsquamous tumors of head and neck, 593
effect of oral contrast agents in dual-modality PET/CT imaging, 784
F-18 FDG PET in diagnosis of breast tumor recurrence and metastases, 250
Posterior tibialis tendon tears, MRI vs. US detection, 51
Prostate imaging, MRI and MRS at 3 Tesla using transceive pelvic phased array coil, 443

R

Rabbit
combined radiofrequency ablation and hot saline injection in rabbit liver, 725
different types of blood pool agents in rabbit MRA-like protocol, 311
dynamic contrast-enhanced MRI of VX2 carcinomas, 539
histologic evolution of high-intensity focused US in muscle, 293
imaging findings after fat graft interposition in injured growth plate, 695
measurement of absolute renal perfusion by MRI, 584
percutaneous radiofrequency thermal ablation of lung tumors, 129
silencing of phosphonate-gadolinium MRI contrast by hydroxyapatite binding, 750
time-related MRI and histopathological changes in radiofrequency ablation lesion in liver, 525
use of additives to improve imaging in plain film and CT urography, 17
vascular stent as active component for locally enhanced MRI, 147
Radiofrequency ablation
combined with hot saline injection in rabbit liver, 725
in vivo efficiency of four ablation systems in pig liver, 609
percutaneous radiofrequency thermal ablation of lung tumors (rabbit model), 129
time-related MRI and histopathological changes in radiofrequency ablation lesion in rabbit liver, 525
Radiography
amorphous silicon, flat-panel, x-ray detector vs. storage phosphor-based computed radiography, 212
chest, monitor luminescence and observer performance for detection of abnormalities, 57
detectability of high- and low-contrast details on a TFT screen and a CRT screen designed for radiologic diagnosis, 719
digital image composition in long-leg radiography with flat-panel detector, 189
femoral anteversion measured by biplane radiography vs. CT, 221
lossy data compression on chest radiography, 243
monitor luminescence and observer performance for detection of abnormalities on chest radiographs, 57
skeletal radiography using flat-panel detector, image quality and clinical acceptance at 50% dose reduction, 230
use of additives to contrast media to improve imaging in plain film and CT urography (rabbit and dog models), 17
Rat
antithrombotic effects of ioxaglate and iohexol and interaction with clopidogrel, 34
helium3 spin-exchange polarization applied to ventilation/perfusion imaging, 334
MRI evaluation of superparamagnetic iron oxide nanoparticles in myocardial infarction, 141
peripheral quantitative CT for detection of diabetic osteopathy, 171
Resistive index, in complete unilateral ureteral obstruction (pig model), 153
RingTag for myocardial centerline assessment, 669

S

Salivary gland tumors, diagnostic accuracy of FDG-PET in nonsquamous tumors of head and neck, 593
Scintigraphy, enalaprilat renal scintigraphy to differentiate cyclosporine toxicity from acute tubular necrosis, 473
Software for coronary artery calcium scoring, diagnostic accuracy, 761
Spinal cord, MR transceive phased array designed for spinal cord imaging at 3 Tesla, 428
Stents
accuracy of computer-aided measurements in stent-graft planning, 164
as active component for locally enhanced MRI (rabbit model), 147
balloon expandable, resistance to hoop stress, 65
carotid artery responses to various self-expanding stents (dog model), 95
coronary artery stent visualization and in-stent stenosis detection using 16-slice CT, 790

T

Tendon tears, posterior tibialis, MRI vs. US detection, 51
Thermal ablation
combined radiofrequency ablation and hot saline injection in rabbit liver, 725
dynamic gadolinium uptake in thermally treated canine brain, 102
histologic evolution of high-intensity focused US in rabbit muscle, 293
in vivo efficiency of four radiofrequency ablation systems in pig liver, 609
percutaneous radiofrequency thermal ablation of lung tumors (rabbit model), 129
time-related MRI and histopathological changes in radiofrequency ablation lesion in rabbit liver, 525
Thrombosis, antithrombotic effects of ioxaglate and iohexol and interaction with clopidogrel (rat model), 34
Thyroid tumors, diagnostic accuracy of FDG-PET in nonsquamous tumors of head and neck, 593
Total hip replacement, algorithm for metal artifact reduction in CT, 769
Tumor necrosis factor-α, cytokine profiles after nonionic dimeric contrast medium injection, 776
Tumors
blood oxygenation level-dependent MRV of brain tumors at 3 Tesla, 409

diagnostic accuracy of FDG-PET in nonsquamous tumors of head and neck, 593
dynamic contrast-enhanced MRI of VX2 carcinomas in rabbits, 539
dynamic gadolinium uptake in thermally treated canine brain, 102
F-18 FDG PET in diagnosis of breast tumor recurrence and metastases, 250
optimal timing of helical CT arterial phase imaging in hypervascular hepatocellular carcinoma, 497
percutaneous radiofrequency thermal ablation of lung tumors (rabbit model), 129

U

Ultrasonography (US). *See also* Echocardiography
adaptive image processing technique improves image quality, 257
histologic evolution of high-intensity focused US in rabbit muscle, 293
vs. MRI to detect posterior tibialis tendon tears, 51
multisite quantitative US device in normal and osteoporotic postmenopausal women, 207
resistive index in complete unilateral ureteral obstruction (pig model), 153
sentinel node detection with Doppler US lymphography (dog model), 358
tumor vascularization studies with dynamic MRI and Doppler US (mouse model), 516
Ureteral obstruction, resistive index in (pig model), 153
Urography
gadolinium-enhanced MR fluoroscopy used as micturating cystourethrography, 617
plain film and CT, use of additives to contrast media to improve imaging (rabbit and dog models), 17
USPIO, different types of blood pool agents in rabbit MRA-like protocol, 311

V

Vascular studies
accuracy of computer-aided measurements in stent-graft planning, 164
calcium-dependent endothelial cell injury induced by contrast media, 366
calibration phantoms and scoring thresholds for quantification of coronary calcium, 559
carotid artery responses to various self-expanding stents (dog model), 95
coronary artery stent visualization and in-stent stenosis detection using 16-slice CT, 790
coronary-artery visualization by electron beam tomography vs. multislice spiral CT, 119
detecting small coronary artery calcium lesions using electron beam tomography, 183
diagnostic accuracy of software for coronary artery calcium scoring, 761
dose of contrast agent in assessing coronary stenosis by myocardial contrast echocardiography, 550
effect on neointimal proliferation and restenosis after coronary stenting in pigs, 743
embolization with autologous fibroblast-attached platinum coils in canine carotid artery aneurysms, 281
Gd-BOPTA–enhanced MRA of pelvic arteries, 504
high-resolution MRI of atherosclerotic lesions at 2 Tesla (mouse model), 532
multifunctional coil catheter for gastrointestinal intervention (dog model), 796
quantification of blood flow in small vessels with velocity encoded MRI, 567
reconstruction windows for multislice CT to quantify coronary calcium, 108 (erratum, 241)
resistance to hoop stress in balloon expandable stents, 65
tumor vascularization studies with dynamic MRI and Doppler US (mouse model), 516
vascular stent as active component for locally enhanced MRI (rabbit model), 147
Venography, blood oxygenation level-dependent MRV of brain tumors at 3 Tesla, 409
Ventilation
distribution in lung transplant recipients evaluated by helium3 MRI, 341
helium3 spin-exchange polarization applied to ventilation/perfusion imaging (rat model), 334
VX2 carcinoma, dynamic contrast-enhanced MRI of (rabbit model), 539